



## Corrigendum

## Corrigendum to “Statistical summability and approximation by de la Vallée-Poussin mean” (M. Mursaleen, A. Alotaibi), [Appl. Math. Lett. 24 (2011) 320–324]

M. Mursaleen<sup>a,\*</sup>, Abdullah Alotaibi<sup>b,1</sup><sup>a</sup> Department of Mathematics, Aligarh Muslim University, Aligarh 202002, India<sup>b</sup> Department of Mathematics, King Abdulaziz University, Jeddah 21589, Saudi Arabia

## ARTICLE INFO

## Article history:

Received 3 August 2011

Accepted 14 September 2011

In the paper by M. Mursaleen and A. Alotaibi [Statistical summability and approximation by de la Vallée-Poussin mean, Applied Mathematics Letters, 24 (2011) 320–324], a correction is required on Page 324, lines 13–14:

Now replacing  $T_n(t, x)$  by  $B_k(t, x) = \frac{1}{\lambda_k} \sum_{n \in I_k} T_n(t, x)$ , and for  $\epsilon' > 0$ , write

$$D = \left\{ k \in \mathbb{N} : \|B_k(1, x) - 1\|_\infty + \|B_k(t, x) - x\|_\infty + \|B_k(t^2, x) - x^2\|_\infty \geq \frac{\epsilon'}{K} \right\}.$$

It should be as follows.

Now replacing  $T_n(., x)$  by  $B_k(., x) = \frac{1}{\lambda_k} \sum_{n \in I_k} T_n(., x)$ , and for  $\epsilon' > 0$ , write

$$D = \left\{ k \in \mathbb{N} : \|B_k(f, x) - f(x)\|_\infty \geq \frac{\epsilon'}{K} \right\}.$$

DOI of original article: [10.1016/j.aml.2010.10.014](https://doi.org/10.1016/j.aml.2010.10.014).

\* Corresponding author. Tel.: +91 9411491600.

E-mail addresses: [mursaleenm@gmail.com](mailto:mursaleenm@gmail.com) (M. Mursaleen), [aalotaibi@kau.edu.sa](mailto:aalotaibi@kau.edu.sa) (A. Alotaibi).

<sup>1</sup> Tel.: +966 500354545.